



## King's Research Portal

DOI:

[10.2337/dbi17-0024](https://doi.org/10.2337/dbi17-0024)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Chapple, S., & Mann, G. E. (2017). New Tricks for Nrf2: Therapeutic Targeting to Restore BK-1 Expression? *Diabetes*, 66, 2538-2540. <https://doi.org/10.2337/dbi17-0024>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## New tricks for Nrf2: therapeutic targeting to restore BK-β1 subunit expression?

Sarah J. Chapple\* and Giovanni E. Mann

Cardiovascular Division, King's BHF Centre for Research Excellence, Faculty of Life Sciences & Medicine, 150 Stamford, London SE1 9NH, U.K.

\*Corresponding author: Sarah J. Chapple

Email [sarah.2.chapple@kcl.ac.uk](mailto:sarah.2.chapple@kcl.ac.uk) Tel +44(0)207 848 4830

The transcription factor NF-E2 related factor 2 (Nrf2) is ubiquitously expressed and a master regulator of antioxidant, phase II and proteostatic genes induced by oxidative or electrophilic stress (1; 2). Basally, Nrf2 is continually synthesized and associates with adaptor protein Kelch-like ECH Associated Protein 1 (Keap1), resulting in its ubiquitination and rapid degradation via the 26S proteasome (3-5). Following oxidative challenge, modification of key cysteine residues on Keap1 prevent Nrf2 degradation, allowing *de novo* Nrf2 to translocate to the nucleus and initiate transcription of target genes containing an antioxidant or electrophilic response element (ARE/EpRE) in their promoter region. In cells, failure to activate Nrf2 defenses increases their susceptibility to oxidative damage and dysfunction, contributing to the pathogenesis of various cardiometabolic disease states including hypertension, cardiomyopathy, gestational and/or Type II diabetes (6-8).

In this issue of *Diabetes* (9), Lu *et al.* demonstrate a novel role for Nrf2 in the regulation of the large conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  (BK) channel-β1 subunit. As with other recently identified Nrf2-regulated solute transporters, this channel has no intrinsic antioxidant activity but is nevertheless important in homeostatic control of cell function. BK channels are expressed on vascular endothelial and smooth muscle cells (SMCs) and consist of a tetramer of α subunits, containing a channel pore domain, voltage sensor and  $\text{Ca}^{2+}$ -sensing region. These α subunits are each associated with a tissue-specific modulatory β subunit (isoforms β1-4), altering channel activity (10). The β1 subunit is abundantly expressed in vascular SMCs and enhances the BK channels sensitivity to  $\text{Ca}^{2+}$  (11). Activation of BK channels leads to  $\text{K}^+$  efflux, resulting in membrane hyperpolarization and inhibition of  $\text{Ca}^{2+}$  influx through large conductance L-type  $\text{Ca}^{2+}$  channels, allowing BK channels to sense and modulate intracellular  $\text{Ca}^{2+}$  levels.

In the current study, Lu *et al.* (9) used a mouse model of high fat diet (HFD) to induce hypertension and insulin resistance. They showed that in HFD mice, diminished Nrf2 expression correlates with a 57% reduction in BK-β1 subunit protein expression. The authors also report that reduced BK-β1 subunit expression coincides with reduced BK channel excitability in patch-clamped single coronary artery SMCs treated with the BK-β1 activator DHS-1. Raised mean arterial pressure (MAP) (HFD  $105.2 \pm 3.7$  vs. lean  $91.9 \pm 4.8$  mmHg) and impaired BK-β1 dependent vasodilation of coronary arteries reported by Lu *et al.* are consistent with previous studies supporting a role for BK-β1 in mediating vasodilation and contributing towards the regulation of myogenic tone and sympathetically mediated MAP.

Previously, deletion of the BK- $\beta$ 1 subunit in mice was shown to uncouple transient  $\text{Ca}^{2+}$  release from intracellular stores ( $\text{Ca}^{2+}$  sparks) to BK channel activation, raising MAP (12; 13) with primarily night-time MAP corresponding to peak activity and enhanced sympathetic outflow affected (14). Diminished BK- $\beta$ 1 expression and  $\text{Ca}^{2+}$  sensitivity are also observed in rat models of genetic hypertension (WKY and SHR) (15) and angiotensin II induced hypertension (11). In humans, few studies have explored the contribution of BK- $\beta$ 1 in vascular function, but do indicate BK- $\beta$ 1 may be a useful therapeutic target for cardiometabolic disease prevention.

Previous studies demonstrate an autoregulatory role for BK channels, with blockade of BK channels using IBTX reducing vasodilation in response to increased intraluminal pressure in isolated human skeletal muscle arterioles (16). A common polymorphism in the gene encoding the BK- $\beta$ 1 subunit (E65K genotype) results in augmented  $\text{Ca}^{2+}$  sensitivity of BK channels and is associated with a lower incidence of diastolic hypertension in humans (17). Interestingly, the frequency of this polymorphism and its negative association with diastolic hypertension appears greatest in female participants over the age of 55 (18). The underlying mechanism(s) responsible for age and sex-specific differences in BK channel activation are at this stage unclear. Ovariectomy does not alter BK channel expression (19), nor does E65K alter *in vitro* BK channel activation in response to 17- $\beta$  estradiol (18), an established activator of BK channels (20). In aged male rodents, BK channel  $\alpha$  and  $\beta$ 1 subunit expression declines (21) and oxidation of BK- $\alpha$  subunit Cys<sup>911</sup> reduces the sensitivity of BK channels to  $\text{Ca}^{2+}$ , reducing channel activation (22). It is therefore conceivable that a pro-oxidative environment found during aging, or as in the present HFD study cardiometabolic disease, may be needed to fully dissect the contribution of BK- $\beta$ 1 to blood pressure in male and female cohorts.

Highlighting the influence of redox-regulated transcription on BK- $\beta$ 1 expression, the present study by Lu *et al.* demonstrates that Nrf2 attenuates the expression of the muscle specific E3 ubiquitin ligase MuRF1 by inhibiting pro-oxidant transcription factor NF $\kappa$ B. Using pull-down assays, this group has previously characterized a direct interaction between MuRF1 and the BK- $\beta$ 1 subunit, with silencing of MuRF1 or inhibition of proteasomal degradation leading to BK- $\beta$ 1 stabilization and enhanced BK channel activity in a Type 1 diabetic mouse model (23). Whilst direct interaction of MuRF1 with BK- $\beta$ 1 strongly suggests enhanced proteolysis does indeed contribute to reduced BK- $\beta$ 1 expression in diabetes, inhibition of global proteasomal activity to prevent BK- $\beta$ 1 degradation, as used in this study, will produce off-target effects including the stabilisation of Nrf2. Future studies genetically ablating the BK- $\beta$ 1-MuRF1 binding site or pulse-chase analysis of BK- $\beta$ 1 protein degradation may provide more definitive evidence for the functional importance of this interaction.

Excitingly the current study provides strong evidence that induction of Nrf2 can restore BK- $\beta$ 1 expression and activity both *in vitro* and *in vivo*. To establish that Nrf2 regulates BK- $\beta$ 1 expression, the authors transfected cultured SMCs with Nrf2 overexpressing and silencing adenoviruses. They showed clearly that enhanced Nrf2 levels result in increased expression of BK- $\beta$ 1 mRNA and protein and that suppression of Nrf2 expression reduces BK- $\beta$ 1 levels. These findings suggest that Nrf2's effect on BK- $\beta$ 1 levels are at least partly exerted through transcriptional regulation, consistent with the authors' description of a Nrf2 ARE consensus sequence in the BK- $\beta$ 1 promoter. Further studies are required to resolve the relative contribution of NF $\kappa$ B mediated BK- $\beta$ 1 proteolysis versus Nrf2 dependent

transcriptional upregulation of BK- $\beta$ 1. Perhaps the most convincing evidence underpinning a potential therapeutic role of Nrf2 in stabilizing BK- $\beta$ 1 is the restoration of BK- $\beta$ 1 expression following 10-day administration of FDA-approved Nrf2 activator DMF in HFD treated mice. Acute treatment of human SMCs with DMF confirms upregulation of BK- $\beta$ 1 mRNA and protein, providing further evidence that BK- $\beta$ 1 is a newly identified Nrf2 target gene (see Fig. 1).

Whilst Nrf2 is capable of inducing BK- $\beta$ 1 subunit expression, the observation that basal BK- $\beta$ 1 subunit mRNA expression is stable despite lower Nrf2 protein expression suggests other transcriptional regulators regulate BK- $\beta$ 1 expression and strongly warrant further investigation. Moreover, as both Nrf2 and BK- $\beta$ 1 expression and activity appear to be sensitive to both reactive oxygen species and O<sub>2</sub> tension (22; 24; 25), conducting further *in vitro* studies under a physiological O<sub>2</sub> levels may provide valuable insights into BK channel modulation in physiological and pathological settings, allowing age and sex associated differences in BK activity to be explored and the efficacy of BK- $\beta$ 1 induction to treat various vasculopathies assessed.

### **Acknowledgements**

S.J.C researched and wrote the article, G.E.M contributed to the discussion and reviewed/edited the article. We gratefully acknowledge research support from the British Heart Foundation (FS/13/66/30445, PG/13/1/29801) and a Novel and Emerging Technologies Grant from Heart Research UK (NET01/13).

### **Duality of Interest**

No conflicts of interest relevant to this article are reported.

## References

1. Itoh K, Chiba T, Takahashi S, Ishii T, Igarashi K, Katoh Y, Oyake T, Hayashi N, Satoh K, Hatayama I, Yamamoto M, Nabeshima Y: An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochemical and biophysical research communications* 1997;236:313-322
2. Chapple SJ, Siow RC, Mann GE: Crosstalk between Nrf2 and the proteasome: therapeutic potential of Nrf2 inducers in vascular disease and aging. *The international journal of biochemistry & cell biology* 2012;44:1315-1320
3. Itoh K, Wakabayashi N, Katoh Y, Ishii T, Igarashi K, Engel JD, Yamamoto M: Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes & development* 1999;13:76-86
4. Tong KI, Katoh Y, Kusunoki H, Itoh K, Tanaka T, Yamamoto M: Keap1 recruits Neh2 through binding to ETGE and DLG motifs: characterization of the two-site molecular recognition model. *Molecular and cellular biology* 2006;26:2887-2900
5. McMahon M, Thomas N, Itoh K, Yamamoto M, Hayes JD: Dimerization of substrate adaptors can facilitate cullin-mediated ubiquitylation of proteins by a "tethering" mechanism: a two-site interaction model for the Nrf2-Keap1 complex. *The Journal of biological chemistry* 2006;281:24756-24768
6. Cheng X, Chapple SJ, Patel B, Puszyk W, Sugden D, Yin X, Mayr M, Siow RC, Mann GE: Gestational diabetes mellitus impairs Nrf2-mediated adaptive antioxidant defenses and redox signaling in fetal endothelial cells in utero. *Diabetes* 2013;62:4088-4097
7. Bai Y, Cui W, Xin Y, Miao X, Barati MT, Zhang C, Chen Q, Tan Y, Cui T, Zheng Y, Cai L: Prevention by sulforaphane of diabetic cardiomyopathy is associated with up-regulation of Nrf2 expression and transcription activation. *Journal of molecular and cellular cardiology* 2013;57:82-95
8. Wang Y, Zhang Z, Sun W, Tan Y, Liu Y, Zheng Y, Liu Q, Cai L, Sun J: Sulforaphane attenuation of type 2 diabetes-induced aortic damage was associated with the upregulation of Nrf2 expression and function. *Oxidative medicine and cellular longevity* 2014;2014:123963
9. Lu T, Sun X, Li Y, Chai Q, Wang XL, Lee HC: Role of Nrf2 Signaling in the Regulation of Vascular BK Channel Beta-1 Subunit Expression and BK Channel Function in High Fat Diet-Induced Diabetic Mice. *Diabetes* 2017;
10. Lee US, Cui J: BK channel activation: structural and functional insights. *Trends in neurosciences* 2010;33:415-423
11. Amberg GC, Bonev AD, Rossow CF, Nelson MT, Santana LF: Modulation of the molecular composition of large conductance, Ca(2+) activated K(+) channels in vascular smooth muscle during hypertension. *The Journal of clinical investigation* 2003;112:717-724
12. Brenner R, Perez GJ, Bonev AD, Eckman DM, Kosek JC, Wiler SW, Patterson AJ, Nelson MT, Aldrich RW: Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature* 2000;407:870-876
13. Pluger S, Faulhaber J, Furstenau M, Lohn M, Waldschutz R, Gollasch M, Haller H, Luft FC, Ehmke H, Pongs O: Mice with disrupted BK channel beta1 subunit gene feature abnormal Ca(2+) spark/STOC coupling and elevated blood pressure. *Circulation research* 2000;87:E53-60
14. Xu H, Garver H, Galligan JJ, Fink GD: Large-conductance Ca2+-activated K+ channel beta1-subunit knockout mice are not hypertensive. *American journal of physiology Heart and circulatory physiology* 2011;300:H476-485
15. Amberg GC, Santana LF: Downregulation of the BK channel beta1 subunit in genetic hypertension. *Circulation research* 2003;93:965-971
16. Feng J, Liu Y, Khabbaz KR, Sodha NR, Osipov RM, Hagberg R, Alper SL, Sellke FW: Large conductance calcium-activated potassium channels contribute to the reduced myogenic tone of peripheral microvasculature after cardiopulmonary bypass. *The Journal of surgical research* 2009;157:123-128

17. Fernandez-Fernandez JM, Tomas M, Vazquez E, Orio P, Latorre R, Senti M, Marrugat J, Valverde MA: Gain-of-function mutation in the KCNMB1 potassium channel subunit is associated with low prevalence of diastolic hypertension. *The Journal of clinical investigation* 2004;113:1032-1039
18. Senti M, Fernandez-Fernandez JM, Tomas M, Vazquez E, Elosua R, Marrugat J, Valverde MA: Protective effect of the KCNMB1 E65K genetic polymorphism against diastolic hypertension in aging women and its relevance to cardiovascular risk. *Circulation research* 2005;97:1360-1365
19. Tsang SY, Yao X, Wong CM, Chan FL, Chen ZY, Huang Y: Differential regulation of K<sup>+</sup> and Ca<sup>2+</sup> channel gene expression by chronic treatment with estrogen and tamoxifen in rat aorta. *European journal of pharmacology* 2004;483:155-162
20. Valverde MA, Rojas P, Amigo J, Cosmelli D, Orio P, Bahamonde MI, Mann GE, Vergara C, Latorre R: Acute activation of Maxi-K channels (hSlo) by estradiol binding to the beta subunit. *Science* 1999;285:1929-1931
21. Shi L, Liu X, Li N, Liu B, Liu Y: Aging decreases the contribution of MaxiK channel in regulating vascular tone in mesenteric artery by unparallel downregulation of alpha- and beta1-subunit expression. *Mechanisms of ageing and development* 2013;134:416-425
22. Tang XD, Garcia ML, Heinemann SH, Hoshi T: Reactive oxygen species impair Slo1 BK channel function by altering cysteine-mediated calcium sensing. *Nature structural & molecular biology* 2004;11:171-178
23. Yi F, Wang H, Chai Q, Wang X, Shen WK, Willis MS, Lee HC, Lu T: Regulation of large conductance Ca<sup>2+</sup>-activated K<sup>+</sup> (BK) channel beta1 subunit expression by muscle RING finger protein 1 in diabetic vessels. *The Journal of biological chemistry* 2014;289:10853-10864
24. Resnik E, Herron J, Fu R, Ivy DD, Cornfield DN: Oxygen tension modulates the expression of pulmonary vascular BKCa channel alpha- and beta-subunits. *American journal of physiology Lung cellular and molecular physiology* 2006;290:L761-L768
25. Chapple SJ, Keeley TP, Mastronicola D, Arno M, Vizcay-Barrena G, Fleck R, Siow RC, Mann GE: Bach1 differentially regulates distinct Nrf2-dependent genes in human venous and coronary artery endothelial cells adapted to physiological oxygen levels. *Free radical biology & medicine* 2016;92:152-162

## Figure Legend

### **Fig. 1 Proposed transcriptional and non-transcription regulation of BK-β1 by Nrf2**

**1.** Large conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  (BK) channels become activated in response to a rise in intracellular  $\text{Ca}^{2+}$ , with the  $\beta 1$  subunit (expressed in smooth muscle cells) increasing channel sensitivity to  $\text{Ca}^{2+}$ .  $\text{K}^+$  extrusion leads to membrane hyperpolarization, promoting inhibition of voltage dependent  $\text{Ca}^{2+}$  channels (VDCC) such as large conductance L-type  $\text{Ca}^{2+}$  channels, thereby preventing contraction. BK  $\alpha$  and  $\beta$  subunit expression and/or activity can be further modulated by aging, oxidative stress induced BK- $\alpha$  Cys<sup>911</sup> oxidation or  $\text{O}_2$  tension. **2.** DMF relieves Keap1-mediated repression of Nrf2, allowing Nrf2 nuclear accumulation and induction of newly identified target gene BK- $\beta 1$ . **3.** Nrf2 also attenuates NF $\kappa$ B stabilization, reducing MuRF1 expression and consequently ubiquitination and degradation of BK- $\beta 1$  via the 26S proteasome.